Covariant Non-local Chiral Quark Model and Pion-photon Transition Distribution Amplitudes

Piotr Kotko* and Michal Praszalowicz†
M. Smoluchowski Institute of Physics, Jagiellonian University, ul. Reymonta 4, 30-059 Kraków, Poland

Using non-local chiral quark model with simple pole ansatz for mass dependence on momentum and non-local currents satisfying Ward-Takahashi identities, we calculate pion to photon transition distribution amplitudes and relevant form factors. For vector amplitude we recover correct normalization fixed by the axial anomaly. We find that due to the non-locality in the vector current, the value of axial form factor at zero momentum transfer is lowered with respect to the vector form factor. Such behaviour – consistent with experiment – is not seen in the local models. Where possible we compare our results to the experimental data.

arXiv:0907.4044v2 [hep-ph] 30 Jul 2009

I. INTRODUCTION

Transition Distribution Amplitudes (TDA) were firstly introduced in ref. [1] as objects parametrizing soft part of the amplitudes for hadron-antihadron annihilation $\bar{H}H \to \gamma^*\gamma$ or backward Compton scattering $\gamma^* H \to H\gamma$, where the hard scale is provided by high virtuality of one of the photons. In some sense TDA's are hybrids of the ordinary Distribution Amplitudes (DA) and Generalized Parton Distributions (GPD) (see [2] for a review). However, if one restricts oneself to the mesonic case, they are more similar to GPD's - the difference is that we deal with matrix elements which are non-diagonal not only in momenta but in the physical states as well. Nevertheless, from kinematical point of view, both are almost identical and therefore we can use similar variables as skewedness for example. In practice we consider two kinds of TDA's: vector and axial, depending on nature of the bilocal quark-antiquark operator sandwiched between photon and meson states. For experimental issues of Transition Distribution Amplitudes see ref. [3].

In order to avoid complexity of bound state physics we limit ourselves to pions only. Pions are Goldstone bosons of broken SU(2) chiral symmetry and their properties are to large extent determined by the symmetry (breaking) alone rather than by the complex phenomenon of confinement. After such simplification, we can use the non-local semi-bosonized Nambu-Jona-Lasinio (NJL) model to get an insight into TDA's. On the other hand, since Transition Distribution Amplitudes are related to anomalous diagrams, they can serve as a demanding tester of the non-local models. We shall come back to this point later in this paper.

First estimates of TDA's were made in refs. [4, 5, 6]. The first one [4] was based on general QCD symmetries supported by simple quark model, while the second [6] and the third one [5] made use of Spectral Quark Model (SQM) [7] and Pauli-Villars regulated NJL, respectively. The calculations in SQM respect all QCD symmetries by definition (i.e. Lorentz invariance, Ward identities, anomalies etc.). In Pauli-Villars regulated NJL model the presence of the finite regulator [5] gives normalization for the vector amplitude which is not consistent with axial anomaly. In [8] we calculated TDA's numerically in the non-local chiral quark model, however, since we have used "naive" currents we also broke the normalization condition of vector TDA.

In the present paper we extend our calculations from ref. [8] in twofold way: we use the full non-local vertices and obtain not only numerical but also analytical results for some of the TDA's. As a result we recover correct normalization conditions fixed by axial anomaly and give predictions for axial form factor which is not restricted by anomaly. Our results are here in qualitative agreement with experimental data. Since we revised our old calculations using different method we traced the mistake in the vector amplitude, which however does not change qualitative results.

The paper is organized as follows. In Section II we review the non-local chiral quark model and give the prescription for modified currents. Next, in Section III, we recall the definitions of TDA's and relevant sum rules. Section IV contains our results with special emphasis on the role of the new pieces coming from the non-local parts of the currents. In Section V we investigate the relevant form factors and finally we summarize our results in Section VI. Technical details are given in the Appendices.

---

*Electronic address: kotko@th.if.uj.edu.pl
†Electronic address: michal@if.uj.edu.pl

II. NON-LOCAL COVARIANT CHIRAL QUARK MODEL

In the following we use the non-local semi-bosonized Namu-Jona-Lasinio model. It is based on the quark-pion interaction in the following form [9]

$$S_{\mathsf{Int}} = \int \frac{d^4k\, d^4l}{(2\pi)^8} \bar{\psi}(k) \sqrt{M(k)} U^{\gamma_5}(k-l) \sqrt{M(l)} \psi(l), \tag{1}$$

where $M(k)$ is dynamical quark mass appearing due to the spontaneous chiral symmetry breaking. Meson field U^{γ_5} is given in terms of the pion field as

$$U^{\gamma_5}(x) = \exp\left\{ \frac{i}{F_\pi} \tau^a \pi^a(x) \gamma_5 \right\}, \tag{2}$$

where $F_\pi = 93\,\mathrm{MeV}$ is the pion decay constant.

One defines mass dependence on momentum as

$$M(k) = M\, F^2(k) \tag{3}$$

where form factor $F(k)$ should vanish for $k \to \infty$ and is chosen to satisfy $F(0) = 1$. Expression for $F(k)$ was obtained analytically in Euclidean space from the instanton model of the QCD vacuum and it is highly non trivial [9]. Therefore here we use the Minkowski form proposed in [10]

$$F(k) = \left(\frac{-\Lambda_n^2}{k^2 - \Lambda_n^2 + i\epsilon} \right)^n, \tag{4}$$

which reproduces reasonably well the instanton result when continued to Euclidean space. However, it can be also used directly in the Minkowski space. Integer parameter n defines a family of models and allows to analyze a dependence of our results on the shape of $F(k)$. It was argued in ref. [11] that $F(k)$ should vanish exponentially for large momenta in order to be consistent with OPE. This is another reason for introducing n parameter, which allows to control high momentum behavior. The method of fixing value of Λ_n is described below.

It is a well known fact that due to the mass dependence on momentum the standard vector and axial currents are not conserved and Ward-Takahashi (WT) identities are not satisfied. Although it was argued in ref. [8] that this violation is not very large and almost does not affect shapes of TDA's, it certainly affects overall normalizations. It is therefore necessary to resolve the problem of normalization if one wants to use TDA's calculated in the non-local model for phenomenological estimates.

There are several ways of constructing conserved currents in presence of non-local interactions see refs. [12, 13, 14, 15, 16]. However, it should be recalled at this point that none of them is unique, because current conservation fixes only the longitudinal part of a given vertex while the transverse one has to be modeled. In this paper we use the simplest "minimal" vertices satisfying WT identities. For vector current $j^\mu = \bar{\psi}\gamma^\mu\psi$ we replace γ^μ by the following non-local vertex

$$\Gamma^\mu(k,p) = \gamma^\mu + g^\mu(k,p), \tag{5}$$

where the non-local addition reads

$$g^\mu(k,p) = -\frac{k^\mu + p^\mu}{k^2 - p^2} \left(M(k) - M(p) \right). \tag{6}$$

This form of non-locality does not introduce singularities as required by general prosperities of vector vertices [13].

Axial current $j_5^\mu = \bar{\psi}\gamma^\mu\gamma_5\psi$ is made conserved by replacing $\gamma^\mu\gamma_5$ by

$$\Gamma_5^\mu(k,p) = \gamma^\mu\gamma_5 + g_5^\mu(k,p), \tag{7}$$

with

$$g_5^\mu(k,p) = \frac{p^\mu - k^\mu}{(p-k)^2} \left(M(k) + M(p) \right) \gamma_5. \tag{8}$$

In contrary to the vector vertex, the axial one contains physical singularity corresponding to the pion.

$M = 225\,\mathrm{MeV}$	
$n = 1$	$\Lambda = 1641\,\mathrm{MeV}$
$n = 5$	$\Lambda = 3823\,\mathrm{MeV}$
$M = 350\,\mathrm{MeV}$	
$n = 1$	$\Lambda = 836\,\mathrm{MeV}$
$n = 5$	$\Lambda = 1970\,\mathrm{MeV}$
$M = 400\,\mathrm{MeV}$	
$n = 1$	$\Lambda = 721\,\mathrm{MeV}$
$n = 5$	$\Lambda = 1704\,\mathrm{MeV}$

Table I: Numerical values of the model parameters obtained using Birse-Bowler formula (9) for the pion decay constant F_π.

For a model given by n we fix Λ_n using the Birse-Bowler [15] formula for the pion decay constant F_π:

$$F_\pi^2 = \frac{N_c}{4\pi^2}\int_0^\infty dk_{\mathrm{E}}^2 \frac{M^2\left(k_{\mathrm{E}}\right) - k_{\mathrm{E}}^2 M\left(k_{\mathrm{E}}\right) M'\left(k_{\mathrm{E}}\right) + k_{\mathrm{E}}^4 M'\left(k_{\mathrm{E}}\right)^2}{\left(k_{\mathrm{E}}^2 + M^2\left(k_{\mathrm{E}}\right)\right)^2} \tag{9}$$

with $F_\pi = 93\,\mathrm{MeV}$. Using (4) this expression can be calculated analytically - see the Appendix of [17]. In Table I we present Λ_n for given values of constituent quark mass M and power n. Let us recall at this point that there exists another expressions for F_π, namely the Pagels-Stokar formula [12]. As shown explicitly in *e.g.* [18] it accommodates a non-local pion-quark interaction, however it corresponds to the naive axial current j_5^μ which, as mentioned above, does not exhibit PCAC. On the contrary the Birse-Bowler formula [15] takes into account the non-local axial current (more precisely its derivative, which is determined unambiguously). See ref. [18] for discussion.

It is important to note that Λ_n *does not correspond* to the QCD scale characteristic for the present model. Indeed function $F(k)$ does not change much for different (n, Λ_n) except for the high momentum part. Therefore, as pointed out in [8, 20], the precise definition of the scale can be done only within QCD, while within an effective model one can only estimate the order of magnitude. For the instanton model the characteristic scale is about $600\,\mathrm{MeV}$ [9]. A somewhat more detailed discussion of this issue can be found in ref. [20].

The non-local model with momentum dependent quark mass given by (4) was applied in the past to several low energy quantities (pion DA [10], two pion generalized DA [19], quark and gluon condensates [20]), using however naive vector and axial currents. Later in ref. [18], pion DA with conserved axial current was estimated using PCAC. Recently a set of photon DA's up to twist-4 was obtained using currents satisfying WT identities [17].

Similar models with however different form of $F(k)$ in (3) were also considered in the literature. In ref. [15] exponential form of $F(k)$ was used. In refs. [21, 22, 23, 24, 25, 26] various distribution amplitudes and correlators have been studied in non-local models similar to ours. Pion and kaon DA were calculated also in [27].

III. KINEMATICS AND DEFINITIONS

Relevant kinematics is very close to the one used in GPD formalism. We consider pion with momentum P_1 and real photon carrying momentum P_2. We shall work in the chiral limit, therefore both $P_1^2 = 0$ and $P_2^2 = 0$. We define momentum transfer $q^\mu = P_2^\mu - P_1^\mu$ and momentum transfer squared $t = q^2$. Two light-like directions are defined by null vectors $n = (1, 0, 0, -1)$ and $\tilde{n} = (1, 0, 0, 1)$. Decomposition of any vector v^μ in this basis reads

$$v^\mu = v^+ \frac{\tilde{n}^\mu}{2} + v^- \frac{n^\mu}{2} + v_T^\mu, \tag{10}$$

while the scalar product

$$u \cdot v = \frac{1}{2} u^+ v^- + \frac{1}{2} u^- v^+ - \vec{u}_T \cdot \vec{v}_T, \tag{11}$$

where arrows denote Euclidean two-vectors. Introducing average momentum $p = \frac{1}{2}\left(P_1 + P_2\right)$ we can define so called skewedness variable

$$\xi = -\frac{q^+}{2p^+}. \tag{12}$$

For massless pions $-1 < \xi < 1$. Switching to the frame where the average momentum does not have transverse part, *i.e.*

$$p^{\mu} = p^{+}\frac{\tilde{n}^{\mu}}{2} + \frac{p^{2}}{p^{+}}\frac{n^{\mu}}{2} \tag{13}$$

we can write the following parametrizations of the pion and photon momenta

$$P_1^{\mu} = (1+\xi)\, p^{+}\frac{\tilde{n}^{\mu}}{2} + (1-\xi)\,\frac{p^{2}}{p^{+}}\frac{n^{\mu}}{2} - \frac{1}{2}q_T^{\mu} \tag{14}$$

$$P_2^{\mu} = (1-\xi)\, p^{+}\frac{\tilde{n}^{\mu}}{2} + (1+\xi)\,\frac{p^{2}}{p^{+}}\frac{n^{\mu}}{2} + \frac{1}{2}q_T^{\mu}. \tag{15}$$

Notice that $p^2 = -t/4$. We denote photon polarization vector by ε. It satisfies condition $\varepsilon \cdot P_2 = 0$.

We can now define TDA's. Vector Transition Distribution Amplitude (VTDA) $V\left(X,\xi,t\right)$ is defined as

$$\int \frac{d\lambda}{2\pi} e^{i\lambda X p^{+}} \left\langle \gamma\left(P_2,\varepsilon\right) \left| \overline{d}\left(-\frac{\lambda}{2}n\right)\Gamma^{\mu} u\left(\frac{\lambda}{2}n\right)\right| \pi^{+}\left(P_1\right)\right\rangle = \frac{i^2 e}{2\sqrt{2}F_{\pi}p^{+}}\varepsilon^{\mu\nu\alpha\beta}\varepsilon^{*}_{\nu}p_{\alpha}q_{\beta}V\left(X,\xi,t\right). \tag{16}$$

In the axial channel the Axial Transition Distribution Amplitude (ATDA) $A\left(X,\xi,t\right)$ is defined as:

$$\begin{aligned}\int \frac{d\lambda}{2\pi} e^{i\lambda X p^{+}} \left\langle \gamma\left(P_2,\varepsilon\right) \left| \overline{d}\left(-\frac{\lambda}{2}n\right)\Gamma_5^{\mu} u\left(\frac{\lambda}{2}n\right)\right| \pi^{+}\left(P_1\right)\right\rangle = {}& \frac{ie}{2\sqrt{2}F_{\pi}p^{+}}P_2^{\mu} q\cdot\varepsilon^{*} A\left(X,\xi,t\right) \\ & + q^{\mu} q\cdot\varepsilon^{*}\frac{ie2\sqrt{2}F_{\pi}\mathrm{sign}\left(\xi\right)}{t}\phi_{\pi}\left(\frac{X+\xi}{2\xi}\right) + \ldots.\end{aligned} \tag{17}$$

Notice that we have explicitly written down the contribution containing massless pole $1/t$ coming from the fact that the pion can couple directly to the axial current. This part is connected with the pion DA $\phi_{\pi}\left(u\right)$. Dots stand for the remaining parts which vanish when contracted with n^{μ} [4].

There exist the following sum rules for TDA's

$$\int_{-1}^{1} dX \left\{ \begin{array}{c} V\left(X,\xi,t\right) \\ A\left(X,\xi,t\right) \end{array} \right. = \frac{2\sqrt{2}F_{\pi}}{m_{\pi}} \left\{ \begin{array}{c} F_V\left(t\right) \\ F_A\left(t\right) \end{array} \right. = 2\sqrt{2}F_{\pi} \left\{ \begin{array}{c} F_V^{\chi}\left(t\right) \\ F_A^{\chi}\left(t\right) \end{array} \right. , \tag{18}$$

where $F_V\left(t\right)$ and $F_A\left(t\right)$ are vector and axial form factors respectively. Quantity $F^{\chi}_{V,A}\left(t\right) = F_{V,A}\left(t\right)/m_{\pi}$ is introduced because we work in the chiral limit (m_{π} is the mass of the pion). Moreover the transition form factor for the process $\pi^0 \to \gamma^{*}\gamma$ is related to the vector form factor by the formula

$$F_{\pi\gamma}\left(t\right) = \sqrt{2}F_V^{\chi}\left(t\right). \tag{19}$$

The value of $F_{\pi\gamma}$ for $t = 0$ is fixed by the axial anomaly and equals

$$F_{\pi\gamma}\left(0\right) = \frac{1}{4\pi^2 F_{\pi}} \approx 0.272\,\mathrm{GeV}^{-1}. \tag{20}$$

There is no such a constraint for the axial form factor. See section V for the discussion.

IV. TDA'S IN COVARIANT NON-LOCAL MODEL

In this section we outline some points of the calculations within the model introduced in the previous Section and show our results. More details are given in the Appendix A and B.

A. Vector TDA

In the vector channel direct calculation of the matrix elements gives

$$\int \frac{d\lambda}{2\pi} e^{i\lambda X p^{+}} \left\langle \gamma\left(P_2,\varepsilon\right) \left| \overline{d}\left(-\frac{\lambda}{2}n\right) \Gamma^{\mu}\, u\left(\frac{\lambda}{2}n\right) \right| \pi^{+}\left(P_1\right) \right\rangle$$

$$= -\frac{\sqrt{2}eMN_c}{F_\pi}\left(Q_d \mathcal{M}_V^{\mu\nu}\left(X,\xi,t\right) + Q_u \mathcal{M}_V^{\mu\nu}\left(-X,\xi,t\right)\right)\varepsilon_\nu^{*}, \quad (21)$$

where N_c is number of colors, Q_u, Q_d are charges of the pertinent quarks and

$$\mathcal{M}_V^{\mu\nu} = \int \frac{d^4k}{\left(2\pi\right)^4}\,\delta\left(k^{+} - \left(X-1\right)p^{+}\right) \times$$

$$\mathrm{Tr}\left\{S\left(k+P_2\right)\Gamma^{\mu}\left(k+P_2,k+P_1\right)S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)\Gamma^{\nu}\left(k+P_1,k\right)\right\}$$

with

$$S\left(p\right) = \frac{1}{\not{p} - M\left(p\right)}. \quad (22)$$

In the formula above quark-pion coupling reads

$$\Gamma_5\left(k,p\right) = F\left(k\right)\gamma_5 F\left(p\right). \quad (23)$$

Notice that we have used the non-local vertex (5) also in the bilocal currents. This is necessary if one wants to maintain WT identities.

Amplitude $\mathcal{M}_V^{\mu\nu}$, due to the natural splitting of the vector vertices to local and non-local parts, can be divided into four pieces

$$\mathcal{M}_V^{\mu\nu} = \mathcal{M}_V^{\mu\nu,\,(0)} + \mathcal{M}_V^{\mu\nu,\,(1)} + \mathcal{M}_V^{\mu\nu,\,(2)} + \mathcal{M}_V^{\mu\nu,\,(3)}. \quad (24)$$

To list them we introduce shorthand notation

$$\hat{dk} = \frac{d^4k}{\left(2\pi\right)^4}\,\delta\left(k^{+} - \left(X-1\right)p^{+}\right). \quad (25)$$

Then

$$\mathcal{M}_V^{\mu\nu,\,(0)} = \int \hat{dk}\, \mathrm{Tr}\left\{S\left(k+P_2\right)\gamma^{\mu}S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)\gamma^{\nu}\right\} \quad (26)$$

corresponds to the quantities calculated in [8]. The new pieces are

$$\mathcal{M}_V^{\mu\nu,\,(1)} = \int \hat{dk}\, \mathrm{Tr}\left\{S\left(k+P_2\right)\gamma^{\mu}S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)g^{\nu}\left(k+P_1,k\right)\right\}, \quad (27)$$

$$\mathcal{M}_V^{\mu\nu,\,(2)} = \int \hat{dk}\, \mathrm{Tr}\left\{S\left(k+P_2\right)g^{\mu}\left(k+P_2,k+P_1\right)S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)\gamma^{\nu}\right\}. \quad (28)$$

The remaining part of $\mathcal{M}_V^{\mu\nu}$ is zero, simply due to vanishing of the Dirac trace, $\mathcal{M}_V^{\mu\nu,\,(3)} = 0$.

Above decomposition, after projecting on the proper tensor structures, leads straightforwardly to the expression for VTDA

$$V\left(X,\xi,t\right) = V^{(0)}\left(X,\xi,t\right) + V^{(1)}\left(X,\xi,t\right) + V^{(2)}\left(X,\xi,t\right). \quad (29)$$

The $V^{(1)} + V^{(2)}$ part is the addition that is required in order to recover correct normalization. The explicit expressions, together with some details needed to calculate $V^{(0)}$ analytically are given in the Appendix A.

Our results are shown in fig. 1. We plot also separately the local part $V^{(0)}$ and the addition $V^{(1)} + V^{(2)}$ itself. We compare the new result with the local version of our model calculated in [8]. Qualitative behavior with M, ξ, n and t is the same as in [8] therefore we do not discuss this further. However now the condition coming from axial anomaly

$$\int_{-1}^{1} V\left(X,\xi,t\right) = \frac{1}{2\pi^2} \quad (30)$$

is satisfied automatically for any value of n and Λ (which does not have to be necessarily equal Λ_n).



Figure 1: Vector TDA for $M = 350\,\mathrm{MeV}$, $t = -0.1\,\mathrm{GeV}^2$ and $n = 1$. a) Solid line representing full VTDA for $\xi = 0.5$ is the sum of the part coming from the local part of the vector vertex (dotted) and the non-local one (dashed); b) comparison of the results of this paper for $\xi = 0.5$ (dashed), $\xi = -0.5$ (dotted) and the local version of the model $M(k) = M$ for $\xi = \pm 0.5$.

B. Axial TDA

For the matrix element of the axial operator we have

$$\int \frac{d\lambda}{2\pi} e^{i\lambda X p^+} \left\langle \gamma\left(P_2,\varepsilon\right) \middle| \overline{d}\left(-\frac{\lambda}{2}n\right) \Gamma_5^\mu\, u\left(\frac{\lambda}{2}n\right) \middle| \pi^+\left(P_1\right) \right\rangle$$
$$= -\frac{\sqrt{2}eMN_c}{F_\pi}\left(Q_d \mathcal{M}_A^{\mu\nu}\left(X,\xi,t\right) - Q_u \mathcal{M}_A^{\mu\nu}\left(-X,\xi,t\right)\right)\varepsilon_\nu^*, \qquad (31)$$

where

$$\mathcal{M}_A^{\mu\nu} = \int \frac{d^4k}{\left(2\pi\right)^4}\,\delta\left(k^+ - \left(X-1\right)p^+\right)\times$$
$$\mathrm{Tr}\left\{S\left(k+P_2\right)\Gamma_5^\mu\left(k+P_2,k+P_1\right)S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)\Gamma^\nu\left(k+P_1,k\right)\right\}.$$

This expression looks very similar to (21), however here full axial vertex Γ_5^μ given in (7) appears instead of the vector one.

Exploring this expression we find that it has again four parts, similarly to the vector case. However only two of them contribute to the ATDA, namely:

$$\mathcal{M}_A^{\mu\nu,\,(0)} = \int \hat{dk}\,\mathrm{Tr}\left\{S\left(k+P_2\right)\gamma^\mu\gamma_5 S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)\gamma^\nu\right\} \qquad (32)$$

and

$$\mathcal{M}_A^{\mu\nu,\,(1)} = \int \hat{dk}\,\mathrm{Tr}\left\{S\left(k+P_2\right)\gamma^\mu\gamma_5 S\left(k+P_1\right)\Gamma_5\left(k+P_1,k\right)S\left(k\right)g^\nu\left(k+P_1,k\right)\right\} \qquad (33)$$

The remaining terms are connected with pion DA (or they are gauge artifacts). This can be seen by noting that

$$g_5^\mu\left(k+P_2,k+P_1\right) = \frac{q^\mu}{t}\left(M\left(k+P_2\right)+M\left(k+P_1\right)\right)\gamma_5 \qquad (34)$$

and comparing with definition (17). Therefore we have

$$A\left(X,\xi,t\right) = A^{(0)}\left(X,\xi,t\right) + A^{(1)}\left(X,\xi,t\right), \qquad (35)$$

where again $A^{(0)}$ is the old part already calculated in [8]. All further details concerning calculations are relegated to Appendix B.

The results are shown in fig. 2. It turns out that the integral over dX of $A^{(1)}\left(X,\xi,t\right)$ part is negative, what shifts the value of the axial form factor towards the experimental value. For negative values of ξ the addition $A^{(1)}\left(X,\xi,t\right)$



Figure 2: Axial TDA for $M = 350\,\mathrm{MeV}$, $t = -0.1\,\mathrm{GeV}^2$ and $n = 1$. a) Solid line representing full ATDA for $\xi = 0.5$ is the sum of the part coming from the local part of the vector vertex (dotted) and the non-local one (dashed). Dash-dotted curve is the results for the local model $M\left(k\right) = M$. b) The same for $\xi = -0.5$; c) results for positive $\xi = 0.5$, $n = 1$, $t = -0.1\,\mathrm{GeV}^2$ and different values of M; d) the same for negative $\xi = -0.5$.

has in a sense unexpected shape, what quite drastically changes the shape of full ATDA for $\xi < 0$. This effect is weaker for lower constituent quark masses as the addition $A^{(1)}$ is proportional to the third power of M/Λ_n. We have checked that for any ξ (both positive and negative)

$$\int_{-1}^{1} dX\, A^{(1)}\left(X, \left|\xi\right|, t\right) = \int_{-1}^{1} dX\, A^{(1)}\left(X, -\left|\xi\right|, t\right) = f\left(t\right) < 0, \tag{36}$$

where f is some function of t only. Equation (36) is a special case of the polynomiality condition, which is satisfied in our model, both for VTDA and ATDA.

V. FORM FACTORS

As already mentioned in Section IV A vector and transition form factors at zero momentum transfer are reproduced correctly, as given by axial anomaly. Full result for transition form factor multiplied by momentum transfer is shown in fig. 3. We make a comparison with CELLO [28], CLEO [29] and some new BaBar data [30] for several values of model parameters. Since our approach is reliable at low momenta transfer we limited ourselves to the region of a few GeV^2. We do not use the standard dipol parametrization for the transition form factor [29] as a reference, because recent BaBar data seem to reveal different behavior (see also [31, 32]). We find that model parameters $M = 300\,\mathrm{MeV}$ and $n = 1$ fits the data quite well, while low constituent quark masses lie much below. However, we underline that momentum transfer around a few GeV^2 can be too large to be treated within the considered approach.

In the case of the axial form factor, we find that due to the contribution of $A^{(1)}$ its value at zero momentum transfer $F_A\left(0\right)$ is lower than for the vector one. The values of $F_A\left(0\right)$ and the relevant ratios $F_A\left(0\right)/F_V\left(0\right)$ for several model



Figure 3: Pion to photon transition form factor times momentum transfer for several model parameters versus CELLO, CLEO and BaBar data in the $0-8\,\mathrm{GeV}^2$ range.

M [MeV]	n	$F_A(0)$	$F_A(0)/F_V(0)$
225	1	0.0217	0.80
350	1	0.0168	0.62
350	5	0.0163	0.60
400	1	0.0161	0.60
400	5	0.0152	0.56

Table II: Numerical values for axial form factor at zero momentum transfer $F_A(0)$ for different model parameters.

parameters are presented in Table II. We notice that when n is increasing, the value of $F_A(0)$ is decreasing very slowly. Therefore we conclude that $F_A(0)$ is quite robust as far as model parameters are concerned. Experimental value given by PDG is

$$F_A^{\exp}(0) = 0.0115 \pm 0.0005 \tag{37}$$

and

$$\left(F_A(0)/F_V(0)\right)_{\exp} = 0.7^{+0.6}_{-0.2}. \tag{38}$$

Although our predictions are much better then the ones obtained in local models (with $F_A(0) = F_V(0)$) [4, 5, 6, 8], it still does not agree with experimental value. Notice however that also prediction for $F_V(0)$ provided by conservation of vector current and axial anomaly

$$F_V(0) \approx 0.027 \tag{39}$$

overshoots the experimental value (PDG)

$$F_V^{\exp}(0) = 0.017 \pm 0.008. \tag{40}$$

We recall also that our calculations were done in the chiral limit. The effects due to the finite pion mass were discussed in [5] within local NJL model and turn out to be small.

VI. SUMMARY

In the present paper w have used the instanton motivated non-local chiral quark model to compute pion-to-photon transition distributions amplitudes in vector and axial channel defined in eqs.(16) and (17). Our approach follows closely the approach of ref. [8] with one important modification. In ref.[8] we have used the momentum dependent constituent quark mass and the nonlocal pion-quark interaction, however, both vector and axial currents were not modified to satisfy Ward-Takahashi identities. We have argued in [8] that such an approach – although inconsistent

– gives reasonable shape for the TDA's missing, however, normalization. Indeed, our present findings confirm this conclusion.

In order to satisfy WT identities the "naive" QCD currents require modifications if the quark mass depends on momentum. Such modifications are not unique because WT identities do not fix the longitudinal part of the current. In the present paper we have used the simplest possible modifications of the vector and axial currents given by eqs.(5) and (7).

More importantly, we have used the nonlocal vertex not only for the photon coupling to the quark loop but also for the vector operator defining the TDA. This modification is not obvious for the following reason. Factorization of the physical process into hard and soft parts is done in QCD in terms of the operator product expansion and the relevant operators entering the definition of the soft part are therefore indeed the "naive" QCD operators. In our language this would correspond to neglecting $\mathcal{M}_V^{\mu\nu,\,(2)}$ of eq. (28). We have checked that this would lead to the violation of the normalization condition for VTDA. In the case of ATDA the non-local part of the axial current does not contribute, since it enters the piece proportional to the pion DA – see eq.(17) – but not the ATDA itself.

Our findings can be summarized as follows. The normalization of TDA's in the vector channel – which is fixed by the axial anomaly – turns out to be correctly reproduced in our approach. At the same time normalization of ATDA's is lowered due to the non-local term in the vector current. As a result the two normalizations are no longer equal, as observed experimentally. The shapes of the TDA's are similar to the ones obtained in the local model and in the non-local model with naive currents. In the case of VTDA's we observe, however, small difference between distributions with positive and negative ξ what was previously not the case. ATDA's acquire negative contribution from the non-local piece which makes the differences between positive and negative ξ's even more profound than in the previously considered models. We have also investigated dependence of TDA's on the choice of constituent quark mass M and the shape of the cutoff function $F(k)$. Dependence on the power n entering $F(k)$ is rather weak, whereas the dependence on M is effectively important only for ATDA for negative ξ.

We have also calculated the corresponding form factors. Axial form factor is quite robust with respect to the shape of the mass dependence on momentum. Pion-to-photon transition form factor, which is directly related to the vector form factor, can be compared with experimental data. We compare it with CELLO, CLEO and recent BaBar data in low energy regime in fig.(3) and conclude that our results reproduce quite well its t dependence.

Let us finish by a remark that we have worked in the chiral limit where $m_\pi = 0$. The effects of adding small current masses are expected to be small, however, only further study can show whether they will be able to correct small deviations from experimental values of $F_{V,A}(0)$. Finally the modifications of the nonlocalities that are still possible as far as longitudinal part of the currents is concerned deserve further investigation.

Acknowledgments

The authors are grateful to W. Broniowski, E. Ruiz Arriola and A. Dorokhov for discussions. The paper was partially supported by the Polish-German cooperation agreement between Polish Academy of Science and DFG.

Appendix A: VECTOR TDA

In the beginning we introduce a shorthand notation in order to make the formulae more compact: following abbreviation

$$M\left(k\right) \equiv M_k \tag{A1}$$

and similarly for other k-dependent quantities. We recall that $M_k = M\,F_k^2$. For the inverse scalar propagators we use

$$D_k = k^2 - M_k^2. \tag{A2}$$

Let us now turn to VTDA. We obtained the following expressions for subsequent pieces giving contribution to $V\left(X,\xi,t\right)$

$$V^{(i)}\left(X,\xi,t\right) = 16i\,MN_c p^+\left(Q_d\mathcal{I}^{(i)}\left(X,\xi,t\right) + Q_u\mathcal{I}^{(i)}\left(-X,\xi,t\right)\right) \tag{A3}$$

where

$$\mathcal{I}^{(0)}\left(X,\xi,t\right) = \int \hat{dk}\,\frac{F_kF_{k+P_1}}{D_{k+P_1}D_{k+P_2}D_k} \times \left\{F_1\left(M_{k+P_1} - M_{k+P_2}\right) + F_2\left(M_k - M_{k+P_2}\right) + M_k\right\}, \tag{A4}$$

$$\mathcal{I}^{(1)}\left(X,\xi,t\right)=\int\hat{dk}\,\frac{\left(-k_T^2\sin^2\theta_T\right)\,F_kF_{k+P_1}\left(M_k-M_{k+P_2}\right)}{D_{k+P_1}D_{k+P_2}D_k\left(k\cdot P_2\right)},\tag{A5}$$

$$\mathcal{I}^{(2)}\left(X,\xi,t\right)=-\int\hat{dk}\,\frac{\left(-k_T^2\sin^2\theta_T\right)\,F_kF_{k+P_1}\left(M_{k+P_2}-M_{k+P_1}\right)}{D_{k+P_1}D_{k+P_2}D_k\left(k\cdot q\right)}.\tag{A6}$$

In the above formulae

$$F_1=-\frac{\vec{k}_T\cdot\vec{q}_T}{t\left(1+\xi\right)}-\frac{1}{2}\left(X-1\right),\tag{A7}$$

$$F_2=2\xi\frac{\vec{k}_T\cdot\vec{q}_T}{t\left(\xi^2-1\right)}+\left(X-1\right)\tag{A8}$$

and θ_T is an angle between $\vec{k}_T$ and $\vec{q}_T$ in the transverse plane. Integration measure $\hat{dk}$ was introduced in eq. (25).

As an example we briefly explain how to calculate $\mathcal{I}^{(1)}$. Following refs. [10] we introduce dimensionless variables $\kappa=k/\Lambda$, $\bar{P}_1=P_1/\Lambda$, $\bar{P}_2=P_2/\Lambda$, $r=M/\Lambda$ and use

$$u_{1,2}=\left(\kappa+\bar{P}_{1,2}\right)^2-1+i\epsilon,\tag{A9}$$

$$u_3=\kappa^2-1+i\epsilon.\tag{A10}$$

Then using (4) we have

$$\mathcal{I}^{(1)}\left(X,\xi,t\right)=-\frac{r}{\Lambda^2}\int\hat{d\kappa}\,\frac{\kappa_T^2\sin^2\theta_T\,u_1^{3n}u_2^{2n}u_3^{n}\left(u_2^{2n}-u_3^{2n}\right)}{\left(\kappa\cdot\bar{P}_2\right)G\left(u_1\right)G\left(u_2\right)G\left(u_3\right)},\tag{A11}$$

where

$$G\left(z\right)=z^{4n+1}+z^{4n}-r^2.\tag{A12}$$

We can obviously write

$$G\left(z\right)=\prod_{i=1}^{4n+1}\left(z-\eta_i\right),\tag{A13}$$

where η_i-s are the roots of $G\left(z\right)=0$ to be obtained numerically. Their properties were discussed in [10] and [17]. Then using decomposition into proper fractions we obtain

$$\mathcal{I}^{(1)}\left(X,\xi,t\right)=-\frac{r}{\Lambda^2}\sum_{i,j,k}^{4n+1}f_if_jf_k\alpha_{ijk}\tilde{\mathcal{I}}_{ijk}^{(1)}\left(X,\xi,t\right),\tag{A14}$$

where

$$f_i=\prod_{j\neq i}^{4n+1}\frac{1}{\left(\eta_i-\eta_j\right)},\tag{A15}$$

$$\alpha_{ijk}=\eta_i^{3n}\eta_j^{2n}\eta_k^{n}\left(\eta_j^{2n}-\eta_k^{2n}\right)\tag{A16}$$

and

$$\tilde{\mathcal{I}}_{ijk}^{(1)}\left(X,\xi,t\right)=\int\hat{d\kappa}\,\frac{\kappa_T^2\sin^2\theta_T}{\kappa\cdot\bar{P}_2\left(u_1-\eta_i\right)\left(u_2-\eta_j\right)\left(u_3-\eta_k\right)}.\tag{A17}$$

Next we assume that this expression can be continued to the Euclidean space – this is equivalent in Minkowski space to deformation of the $d\kappa^-$ integration contour as described in [10, 17]. This choice of the contour assures that the results are real and analytical in Λ_n. The remaining parts of integrals $\mathcal{I}^{(0)}$, $\mathcal{I}^{(2)}$ can be transformed to a sum of the integrals similarly as in the case $\mathcal{I}^{(1)}$. It is convenient to start with the $d\kappa^-$ integration using residue theorem. Positions of the poles in the κ^- complex plane depend on the kinematical region and on the sign of ξ, however the pole connected with $\kappa \cdot \bar{P}_2$ does not give contribution as it should be. Physical support $-1 < X < 1$ splits into three pieces: $-1 < X < -|\xi|$, $-|\xi| < X < |\xi|$ and $|\xi| < X < 1$. Integrals $\mathcal{I}^{(i)}(X,\xi,t)$ are nonzero in the last two, while $\mathcal{I}^{(i)}(-X,\xi,t)$ in the first two intervals.

In order to write down the results it is useful to introduce the following notation

$$g_i^{(A)} = \left(\kappa_T^2+1\right)(1-\xi) - \vec{\kappa}_T\cdot\vec{q}_T\,(X+1) - (X-1)\left(1-3\xi-X\left(1+\xi\right)\right)\bar{p}^2 + (1-\xi)\,\eta_i, \tag{A18}$$

$$g_i^{(B)} = \left(\kappa_T^2+1\right)(1-\xi) - \vec{\kappa}_T\cdot\vec{q}_T\,(X-1) - (X-1)^2\,(1+\xi)\,\bar{p}^2 + (1-\xi)\,\eta_i, \tag{A19}$$

$$h_i^{(A)} = \left(\kappa_T^2+1\right)\xi + X\,\vec{\kappa}_T\cdot\vec{q}_T - \left(X^2-1\right)\xi\bar{p}^2 + \xi\eta_i, \tag{A20}$$

$$h_i^{(B)} = \left(\kappa_T^2+1\right)\xi + (X-1)\;\vec{\kappa}_T\cdot\vec{q}_T - (X-1)^2\,\xi\bar{p}^2 + \xi\eta_i, \tag{A21}$$

$$d_{ij}^{(AB)} = -2\xi\left(\kappa_T^2+1\right) - 2X\,\vec{\kappa}_T\cdot\vec{q}_T + 2\left(X^2-1\right)\xi\bar{p}^2 + (X-\xi)\,\eta_i - (X+\xi)\,\eta_j, \tag{A22}$$

$$d_{ij}^{(AC)} = -(1+\xi)\left(\kappa_T^2+1\right) - (1-X)\;\vec{\kappa}_T\cdot\vec{q}_T + (1-\xi)\,(X-1)^2\,\bar{p}^2 + (X+\xi)\,\eta_i - (X-1)\,\eta_j, \tag{A23}$$

$$d_{ij}^{(CB)} = (1-\xi)\left(\kappa_T^2+1\right) + (1-X)\;\vec{\kappa}_T\cdot\vec{q}_T + (1+\xi)\,(X-1)^2\,\bar{p}^2 + (X-\xi)\,\eta_i - (X-1)\,\eta_j, \tag{A24}$$

$$\beta_{ijk} = -\alpha_{kji} \tag{A25}$$

where $\bar{p} = p/\Lambda$ is dimensionless average momentum. Then we have

- $|\xi| < X < 1$

$$\mathcal{I}^{(0)}\left(X,|\xi|,t\right) = \mathcal{I}^{(0)}\left(X,-|\xi|,t\right) = \frac{1}{2}\frac{i\,r}{\Lambda p^+}\;(X-1)\sum_{i,j,k}^{4n+1} f_i f_j f_k \int \frac{d^2\kappa_T}{(2\pi)^3}\,\frac{\alpha_{ijk}F_2 + \alpha_{kji}F_1 + \eta_i^{3n}\eta_j^{4n}\eta_k^{n}}{d_{ki}^{(AC)}d_{kj}^{(CB)}} \tag{A26}$$

$$\mathcal{I}^{(1)}\left(X,|\xi|,t\right) = \mathcal{I}^{(1)}\left(X,-|\xi|,t\right) = \frac{i\,r}{\Lambda p^+}\;(X-1)^2\sum_{i,j,k}^{4n+1} \alpha_{ijk} f_i f_j f_k \int \frac{d^2\kappa_T}{(2\pi)^3}\,\frac{-\kappa_T^2\sin^2\theta_T}{g_k^{(B)}d_{ki}^{(AC)}d_{kj}^{(CB)}} \tag{A27}$$

$$\mathcal{I}^{(2)}\left(X,|\xi|,t\right) = \mathcal{I}^{(2)}\left(X,-|\xi|,t\right) = \frac{1}{2}\frac{i\,r}{\Lambda p^+}\;(X-1)^2\sum_{i,j,k}^{4n+1} \beta_{ijk} f_i f_j f_k \int \frac{d^2\kappa_T}{(2\pi)^3}\,\frac{-\kappa_T^2\sin^2\theta_T}{h_k^{(B)}d_{ki}^{(AC)}d_{kj}^{(CB)}} \tag{A28}$$

- $-|\xi| < X < |\xi|$

$$\mathcal{I}^{(0)}\left(X,|\xi|,t\right) = -\frac{1}{2}\frac{i\,r}{\Lambda p^+}\;(X+\xi)\sum_{i,j,k}^{4n+1} f_i f_j f_k \int \frac{d^2\kappa_T}{(2\pi)^3}\,\frac{\alpha_{ijk}F_2 + \alpha_{kji}F_1 + \eta_i^{3n}\eta_j^{4n}\eta_k^{n}}{d_{ij}^{(AB)}\left(-d_{ki}^{(AC)}\right)} \tag{A29}$$

$$\mathcal{I}^{(0)}\left(X,-|\xi|,t\right) = -\frac{1}{2}\frac{i\,r}{\Lambda p^+}\;(X-\xi)\sum_{i,j,k}^{4n+1} f_i f_j f_k \int \frac{d^2\kappa_T}{(2\pi)^3}\,\frac{\alpha_{ijk}F_2 + \alpha_{kji}F_1 + \eta_i^{3n}\eta_j^{4n}\eta_k^{n}}{d_{ij}^{(AB)}d_{kj}^{(CB)}} \tag{A30}$$

$$\mathcal{I}^{(1)}\left(X,\left|\xi\right|,t\right) = -\frac{i\,r}{\Lambda p^{+}}\,\left(X+\xi\right)^{2}\sum_{i,j,k}^{4n+1}\alpha_{ijk}f_{i}f_{j}f_{k}\int\frac{d^{2}\kappa_{T}}{\left(2\pi\right)^{3}}\,\frac{-\kappa_{T}^{2}\sin^{2}\theta_{T}}{g_{i}^{(A)}d_{ij}^{(AB)}\left(-d_{ki}^{(AC)}\right)} \tag{A31}$$

$$\mathcal{I}^{(1)}\left(X,-\left|\xi\right|,t\right) = -\frac{i\,r}{\Lambda p^{+}}\,\left(X-\xi\right)^{2}\sum_{i,j,k}^{4n+1}\alpha_{ijk}f_{i}f_{j}f_{k}\int\frac{d^{2}\kappa_{T}}{\left(2\pi\right)^{3}}\,\frac{-\kappa_{T}^{2}\sin^{2}\theta_{T}}{g_{j}^{(B)}d_{ij}^{(AB)}d_{kj}^{(CB)}} \tag{A32}$$

$$\mathcal{I}^{(2)}\left(X,\left|\xi\right|,t\right) = -\frac{1}{2}\frac{i\,r}{\Lambda p^{+}}\,\left(X+\xi\right)^{2}\sum_{i,j,k}^{4n+1}\beta_{ijk}f_{i}f_{j}f_{k}\int\frac{d^{2}\kappa_{T}}{\left(2\pi\right)^{3}}\,\frac{-\kappa_{T}^{2}\sin^{2}\theta_{T}}{h_{i}^{(A)}d_{ij}^{(AB)}\left(-d_{ki}^{(AC)}\right)} \tag{A33}$$

$$\mathcal{I}^{(2)}\left(X,-\left|\xi\right|,t\right) = -\frac{1}{2}\frac{i\,r}{\Lambda p^{+}}\,\left(X-\xi\right)^{2}\sum_{i,j,k}^{4n+1}\beta_{ijk}f_{i}f_{j}f_{k}\int\frac{d^{2}\kappa_{T}}{\left(2\pi\right)^{3}}\,\frac{-\kappa_{T}^{2}\sin^{2}\theta_{T}}{h_{i}^{(A)}d_{ij}^{(AB)}d_{kj}^{(CB)}} \tag{A34}$$

Integral $\mathcal{I}^{(0)}$ can be calculated completely analytically, however we do not present the result here to make the paper more compact. It should be pointed out that the analytic formulae contain large sums of complicated complex expressions – therefore it is usually more efficient to perform all integrals numerically.

Appendix B: AXIAL TDA

In the axial channel we get

$$A^{(i)}\left(X,\xi,t\right) = -16i\,M\,N_{c}\,p^{+}\left(Q_{d}\mathcal{J}^{(i)}\left(X,\xi,t\right)-Q_{u}\mathcal{J}^{(i)}\left(-X,\xi,t\right)\right), \tag{B1}$$

where

$$\begin{aligned}\mathcal{J}^{(0)}\left(X,\xi,t\right) = \int d\hat{k}\,&\frac{F_{k}F_{k+P_{1}}}{D_{k+P_{1}}D_{k+P_{2}}D_{k}}\\ &\times\Big\{F_{1}\left[M_{k+P_{1}}-M_{k+P_{2}}-2M_{k}\right]-F_{2}\left(M_{k+P_{2}}-M_{k}\right)+M_{k}+2F_{3}\left[M_{k+P_{1}}-M_{k}\right]\Big\},\end{aligned} \tag{B2}$$

$$\begin{aligned}\mathcal{J}^{(1)}\left(X,\xi,t\right) = \int d\hat{k}\,&\frac{F_{k}F_{k+P_{1}}\left(M_{k}-M_{k+P_{2}}\right)}{k\cdot P_{2}D_{k+P_{1}}D_{k+P_{2}}D_{k}}\\ \times\Big\{F_{1}&\left[M_{k+P_{1}}M_{k}-M_{k+P_{2}}M_{k}-2k^{2}-2k\cdot p\right]+F_{3}\big[M_{k+P_{1}}M_{k+P_{2}}-M_{k}M_{k+P_{2}}+M_{k}M_{k+P_{1}}\\ &-k^{2}+P_{1}\cdot P_{2}\big]\Big\}.\end{aligned} \tag{B3}$$

Functions F_1, F_2 are the same as in vector case while

$$F_{3} = \frac{8\xi\left(\vec{k}_{T}\cdot\vec{q}_{T}\right)^{2}+2t\,\vec{k}_{T}\cdot\vec{q}_{T}\left(X-1\right)\left(2\xi^{2}+\xi-1\right)+t\,\left(\xi^{2}-1\right)\left(t\,\left(X-1\right)^{2}\left(1+\xi\right)-4\xi k_{T}^{2}\right)}{2\left(1-\xi\right)\left(1+\xi\right)^{2}t^{2}}. \tag{B4}$$

Calculations proceed quite similarly as before, although they are more involved. However, in the case of $\mathcal{J}^{(1)}$ additional difficulty arises if one wants to obtain result for any n. Let us discuss this further.

Problematic part comes from the subterm of $\mathcal{J}^{(1)}$ containing more then 4 powers of $F(k)$. Using (A9),(A10) this part reads

$$\mathcal{J}_{A}^{(1)}\left(X,\xi,t\right) = \frac{r^{3}}{\Lambda^{2}}\int d\hat{k}\,\frac{u_{1}^{n}u_{2}^{2n}\left(u_{2}^{2n}-u_{1}^{2n}\right)\left(F_{1}+F_{3}\right)}{\kappa\cdot\bar{P}_{2}G\left(u_{1}\right)G\left(u_{2}\right)G\left(u_{3}\right)u_{3}^{n}}. \tag{B5}$$

Notice that there is an additional u_3^n in denominator which gives rise to multiple poles if $n \geq 2$. However if $n = 1$

$$\mathcal{J}^{(1)}_{A,\,n=1}\left(X,\xi,t\right) = \frac{r^3}{\Lambda^2}\sum_{i,j}^{5}\sum_{k}^{6} f_i f_j \tilde{f}_k \int d\hat{k}\, \frac{\left(\eta_i\eta_j^4 - \eta_i^3\eta_j^2\right)\left(F_1 + F_3\right)}{\kappa\cdot\bar{P}_2\left(u_1-\eta_i\right)\left(u_2-\eta_j\right)\left(u_3-\eta_k\right)}, \tag{B6}$$

where

$$\tilde{f}_i = \prod_{j=0,j\neq i}^{5}\frac{1}{\left(\eta_i-\eta_j\right)} \tag{B7}$$

and $\eta_0 = 0$, while $\eta_1,\ldots,\eta_5$ remain solutions of $G = 0$. In order to avoid calculating residues of the multiple poles if $n \geq 2$, we choose to close the contour in such a way that multiple pole does not lie inside. The price we pay is that relevant expressions become more complicated. Moreover, although the pole coming from $\kappa\cdot\bar{P}_2$ does not give contribution to $\mathcal{J}^{(1)}$, it does contribute to $\mathcal{J}_A^{(1)}$ and $\mathcal{J}_B^{(1)}$ separately where $\mathcal{J}^{(1)} = \mathcal{J}_A^{(1)} + \mathcal{J}_B^{(1)}$. This fact has to be taken into account when we decide to choose different contours for $\mathcal{J}_A^{(1)}$ and $\mathcal{J}_B^{(1)}$.

Now we are in a position to write the results. We use the same abbreviations as in the vector case, additionally we define:

$$c_{ijk} = -F_1\left(\beta_{ijk} + 2\eta_i^{3n}\eta_j^{4n}\eta_k^{n}\right) - 2F_3\beta_{ijk} + F_2\alpha_{ijk} + \eta_i^{3n}\eta_j^{4n}\eta_k^{n} \tag{B8}$$

$$w_{ijk}^{(A,B,C)} = F_1 r^2\beta_{ijk} - F_1\beta_{ijk}t^{(A,B,C)} + F_3 r^2\left(\alpha_{ijk}+\beta_{ijk}\right) + F_3\beta_{ijk}s^{(A,B,C)}, \tag{B9}$$

where

$$t^{(A)} = \frac{-\left(1+2\xi\right)\kappa_T^2 + \left(1+\eta_i - \vec{\kappa}_T\cdot\vec{q}_T\right)\left(2X-1\right) + \left(X-1\right)\left(X\left(2\xi-1\right)+1\right)\vec{p}^2}{X+\xi} \tag{B10}$$

$$s^{(A)} = \frac{-\left(1+\xi\right)\kappa_T^2 + \left(1+\eta_i-\vec{\kappa}_T\cdot\vec{q}_T\right)\left(X-1\right) - \left(X-1\right)\Big(\left(X-1\right)^2\left(1-\xi\right) + 2\left(X+\xi\right)\Big)\vec{p}^2}{X+\xi} \tag{B11}$$

$$t^{(B)} = \frac{\left(2\xi-1\right)\kappa_T^2 + \left(1+\eta_j+\vec{\kappa}_T\cdot\vec{q}_T\right)\left(2X-1\right) - \left(X-1\right)\left(\left(2X-1\right)\left(1+\xi\right)+X-\xi\right)\vec{p}^2}{X-\xi} \tag{B12}$$

$$s^{(B)} = \frac{\left(-1+\xi\right)\kappa_T^2 + \left(1+\eta_j+\vec{\kappa}_T\cdot\vec{q}_T\right)\left(X-1\right) - \left(X-1\right)\Big(\left(X-1\right)^2\left(1+\xi\right) + 2\left(X-\xi\right)\Big)\vec{p}^2}{X-\xi} \tag{B13}$$

$$t^{(C)} = \frac{\kappa_T^2 + \left(1+\eta_k\right)\left(2X-1\right)}{X-1} + \left(X-1\right)\vec{p}^2 \tag{B14}$$

$$s^{(C)} = 1 + \eta_k - 2\vec{p}^2. \tag{B15}$$

We recall at this point that $\eta_0 = 0$, therefore although sums below run from unity, notation like "d_{i0}" makes sense.

- $|\xi| < X < 1$

$$\mathcal{J}^{(0)}\left(X,\left|\xi\right|,t\right) = \mathcal{J}^{(0)}\left(X,-\left|\xi\right|,t\right) = \frac{i\,r}{\Lambda p^+}\left(X-1\right)\sum_{i,j,k}^{4n+1} f_i f_j f_k \int\frac{d^2\kappa_T}{\left(2\pi\right)^3}\,\frac{c_{ijk}}{d_{ki}^{(AC)}d_{kj}^{(CB)}} \tag{B16}$$

$$\mathcal{J}_A^{(1)}\left(X,|\xi|,t\right) = \mathcal{J}_A^{(1)}\left(X,-|\xi|,t\right) = -\frac{i\,r^3}{\Lambda p^+}\sum_{i,j,k}^{4n+1} f_i f_j f_k \left(\eta_i^n\eta_j^{4n} - \eta_i^{3n}\eta_j^{2n}\right)\int\frac{d^2\kappa_T}{(2\pi)^3}$$

$$(F_1+F_3)\left\{\frac{(X+\xi)^{n+2}}{g_i^{(A)} d_{ij}^{(AB)}\left(-d_{ki}^{(AC)}\right)\left(-d_{0i}^{(AC)}\right)^n} + \frac{(X-\xi)^{n+2}}{g_j^{(B)} d_{ij}^{(AB)} d_{kj}^{(CB)}\left(-d_{0j}^{(CB)}\right)^n} + \frac{(1-\xi)^{n+2}}{\left(-g_i^{(A)}\right) g_j^{(B)} g_k^{(B)}\left(-g_0^{(B)}\right)^n}\right\} \tag{B17}$$

$$\mathcal{J}_B^{(1)}\left(X,|\xi|,t\right) = \mathcal{J}_B^{(1)}\left(X,-|\xi|,t\right) = \frac{i\,r}{\Lambda p^+}\left(X-1\right)^2\sum_{i,j,k}^{4n+1} f_i f_j f_k \int\frac{d^2\kappa_T}{(2\pi)^3}\frac{w_{ijk}^{(C)}}{g_k^{(B)} d_{ki}^{(AC)} d_{kj}^{(CB)}} \tag{B18}$$

- $-|\xi| < X < |\xi|$

$$\mathcal{J}^{(0)}\left(X,|\xi|,t\right) = -\frac{i\,r}{\Lambda p^+}\left(X+\xi\right)\sum_{i,j,k}^{4n+1} f_i f_j f_k \int\frac{d^2\kappa_T}{(2\pi)^3}\frac{c_{ijk}}{d_{ki}^{(AC)} d_{kj}^{(CB)}} \tag{B19}$$

$$\mathcal{J}^{(0)}\left(X,-|\xi|,t\right) = -\frac{i\,r}{\Lambda p^+}\left(X-\xi\right)\sum_{i,j,k}^{4n+1} f_i f_j f_k \int\frac{d^2\kappa_T}{(2\pi)^3}\frac{c_{ijk}}{d_{ij}^{(AB)} d_{kj}^{(CB)}} \tag{B20}$$

$$\mathcal{J}_A^{(1)}\left(X,|\xi|,t\right) = -\frac{i\,r^3}{\Lambda p^+}\sum_{i,j,k}^{4n+1} f_i f_j f_k \left(\eta_i^n\eta_j^{4n} - \eta_i^{3n}\eta_j^{2n}\right)\int\frac{d^2\kappa_T}{(2\pi)^3}\frac{(F_1+F_3)\left(X+\xi\right)^{n+2}}{g_i^{(A)} d_{ij}^{(AB)}\left(-d_{ki}^{(AC)}\right)\left(-d_{0i}^{(AC)}\right)^n} \tag{B21}$$

$$\mathcal{J}_A^{(1)}\left(X,-|\xi|,t\right) = -\frac{i\,r^3}{\Lambda p^+}\sum_{i,j,k}^{4n+1} f_i f_j f_k \left(\eta_i^n\eta_j^{4n} - \eta_i^{3n}\eta_j^{2n}\right)\int\frac{d^2\kappa_T}{(2\pi)^3}\frac{(F_1+F_3)\left(X-\xi\right)^{n+2}}{g_j^{(B)}\left(-d_{ij}^{(AB)}\right)\left(-d_{kj}^{(CB)}\right)\left(-d_{0j}^{(CB)}\right)^n} \tag{B22}$$

$$\mathcal{J}_B^{(1)}\left(X,|\xi|,t\right) = -\frac{i\,r}{\Lambda p^+}\left(X+\xi\right)^2\sum_{i,j,k}^{4n+1} f_i f_j f_k \int\frac{d^2\kappa_T}{(2\pi)^3}\frac{w_{ijk}^{(A)}}{g_i^{(A)} d_{ij}^{(AB)}\left(-d_{ki}^{(AC)}\right)} \tag{B23}$$

$$\mathcal{J}_B^{(1)}\left(X,-|\xi|,t\right) = -\frac{i\,r}{\Lambda p^+}\left(X+\xi\right)^2\sum_{i,j,k}^{4n+1} f_i f_j f_k \int\frac{d^2\kappa_T}{(2\pi)^3}\frac{w_{ijk}^{(B)}}{g_j^{(B)}\left(-d_{ij}^{(AB)}\right)\left(-d_{kj}^{(CB)}\right)} \tag{B24}$$

Notice, that some of the integrals are superficially divergent. Convergence is assured by the identity

$$\sum_{i=1}^{4n+1} f_i\eta_i^N = \begin{cases} 1 & \text{for } N = 4n \\ 0 & \text{for N < 4n} \end{cases}, \tag{B25}$$

see [17] for the proof and discussion.

[1] B. Pire and L. Szymanowski, Phys. Rev. D **71** (2005) 111501 [arXiv:hep-ph/0411387].
[2] A. V. Belitsky and A. V. Radyushkin, Phys. Rept. **418** (2005) 1 [arXiv:hep-ph/0504030].
[3] J. P. Lansberg, B. Pire and L. Szymanowski, Phys. Rev. D **73** (2006) 074014 [arXiv:hep-ph/0602195] and World Scientific (2008), p. 367, arXiv:0709.2567 [hep-ph].
[4] B. C. Tiburzi, Phys. Rev. D **72** (2005) 094001 [arXiv:hep-ph/0508112].

[5] A. Courtoy and S. Noguera, Phys. Rev. D **76** (2007) 094026 [arXiv:0707.3366 [hep-ph]].
[6] W. Broniowski and E. R. Arriola, Phys. Lett. B **649** (2007) 49 [arXiv:hep-ph/0701243].
[7] E. Ruiz Arriola and W. Broniowski, Phys. Rev. D **67** (2003) 074021 [arXiv:hep-ph/0301202].
[8] P. Kotko and M. Praszalowicz, Acta Phys. Polon. B **40** (2009) 123 [arXiv:0803.2847 [hep-ph]].
[9] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **245** (1984) 259 and **272** (1986) 457.
[10] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **64** (2001) 074003 [arXiv:hep-ph/0105188].
[11] A. E. Dorokhov, Eur. Phys. J. C **42** (2005) 309 [arXiv:hep-ph/0505007], JETP Lett. **82**, 1 (2005) [Pisma Zh. Eksp. Teor. Fiz. **82**, 3 (2005)] [arXiv:hep-ph/0505196].
[12] H. Pagels and S. Stokar, Phys. Rev. D **20** (1979) 2947.
[13] J. S. Ball and T. W. Chiu, Phys. Rev. D **22** (1980) 2542.
[14] B. Holdom, J. Terning and K. Verbeek, Phys. Lett. B **232** (1989) 351; Phys. Lett. B **245** (1990) 612.
[15] R. D. Bowler and M. C. Birse, Nucl. Phys. A **582** (1995) 655 [arXiv:hep-ph/9407336]; R. S. Plant and M. C. Birse, Nucl. Phys. A **628** (1998) 607 [arXiv:hep-ph/9705372].
[16] M. R. Frank, K. L. Mitchell, C. D. Roberts and P. C. Tandy, Phys. Lett. B **359** (1995) 17 [arXiv:hep-ph/9412219].
[17] P. Kotko, M. Praszalowicz, *Photon Distribution Amplitudes in Non-local Chiral Quark Model*, in preparation.
[18] A. Bzdak and M. Praszalowicz, Acta Phys. Polon. B **34** (2003) 3401 [arXiv:hep-ph/0305217].
[19] M. Praszalowicz and A. Rostworowski, Acta Phys. Polon. B **34** (2003) 2699 [arXiv:hep-ph/0302269].
[20] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **66** (2002) 054002 [arXiv:hep-ph/0111196].
[21] V. Y. Petrov, M. V. Polyakov, R. Ruskov, C. Weiss and K. Goeke, Phys. Rev. D **59** (1999) 114018 [arXiv:hep-ph/9807229].
[22] A. E. Dorokhov, W. Broniowski and E. Ruiz Arriola, Phys. Rev. D **74** (2006) 054023 [arXiv:hep-ph/0607171].
[23] I. V. Anikin, A. E. Dorokhov and L. Tomio, Phys. Atom. Nucl. **64** (2001) 1329 [Yad. Fiz. **64** (2001) 1405].
[24] A. E. Dorokhov, M. K. Volkov and V. L. Yudichev, Phys. Atom. Nucl. **66** (2003) 941 [Yad. Fiz. **66** (2003) 973] [arXiv:hep-ph/0203136].
[25] A.E. Dorokhov, M.K. Volkov, V.L. Yudichev, JINR-E4-2001-162 (2001)
[26] A. E. Dorokhov and W. Broniowski, Eur. Phys. J. C **32** (2003) 79 [arXiv:hep-ph/0305037].
[27] S. i. Nam, H. C. Kim, A. Hosaka and M. M. Musakhanov, Phys. Rev. D **74** (2006) 014019 [arXiv:hep-ph/0605259]; S. i. Nam and H. C. Kim, Phys. Rev. D **74**, 076005 (2006) [arXiv:hep-ph/0609267] and Phys. Rev. D **74** (2006) 096007 [arXiv:hep-ph/0608018].
[28] H. J. Behrend *et al.* [CELLO Collaboration], Z. Phys. C **49** (1991) 401.
[29] J. Gronberg *et al.* [CLEO collaboration], Phys. Rev. D **57** (1998) 33 (1998).
[30] B. Aubert [The BABAR Collaboration], arXiv:0905.4778 [hep-ex].
[31] A. V. Radyushkin, arXiv:0906.0323 [hep-ph].
[32] M.V. Polyakov, arXiv:0906.0538 [hep-ph].